

Weed	2,4-D	glyphosate	glyphosate + dicamba	glyphosate + 2,4-D	glyphosate + flumioxazin + dicamba	glyphosate + flumioxazin + 2,4-D	glyphosate + rimsulfuron + thifensulfuron	glyphosate + thifensulfuron + tribenuron	glyphosate + saflufenacil	paraquat	paraquat + diuron
Annual bluegrass	N	E	E	E	E	E	E	E	E	GE	E
Little barley	N	E	E	E	E	E	E	E	E	G	E
Buttercups	G	E	E	E	E	E	E	E	E	E	E
Carolina geranium	PF	PF	G	F	GE	F	G	GE	---	GE	E
Chickweed	P	E	E	E	E	E	E	E	E	E	E
Cudweed	NP	E	E	E	E	E	E	E	E	FG	G
Curly dock	F	F	GE	FG	F	FG	F	E	---	NP	P
Cutleaf eveningprimrose	E	PF	G	E	GE	E	PF	F	---	F <sup>1</sup>	G <sup>1</sup>
Field pansy	P	F	F	F	F	F	--	F	---	G	GE
Henbit	PF	G	E	E	E	E	E	E	---	E	E
Horseweed (marestail)	GE <sup>2</sup>	GE <sup>3</sup>	E	E <sup>2</sup>	E	E <sup>2</sup>	GE <sup>3</sup>	GE <sup>3</sup>	E	PF	G
Ryegrass	N	G	G	G	E	G	GE	G	E	F	FG
Speedwell	PF	E	E	E	E	E	E	E	E	E	E
Swinecress	F	FG	FG	G	G	G	---	GE	---	P	G
Vetch	E	F	E	E	E	E	---	G	---	PF	F
Virginia pepperweed	GE	G	GE	E	E	E	G	G	---	G	G
Wheat or rye cover crop	N	E	E	E	E	E	E	E	E	G <sup>4</sup>	GE <sup>4</sup>
Wild mustard, wild radish	FG <sup>5</sup>	FG	G	E	G	E	FG	GE	---	FG	G

Note: E = excellent, 90% or better control; G = good, 80 to 90% control; F = fair, 50 to 80% control; P = poor, 25 to 50% control; N = no control, less than 25% control

<sup>1</sup> This level of control requires that the primrose be blooming when treated.

<sup>2</sup> This level of control requires 2 pt of 2,4-D.

<sup>3</sup> Control will be poor on glyphosate-resistant horseweed.

<sup>4</sup> Wheat or rye must have visible seed heads for this level of control.

<sup>5</sup> Wild radish and wild mustard control by 2,4-D is good if application is made before plants begin flowering. Use 1 pt per acre of 2,4-D to control these species.